

PCT

WIPO

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference ?	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/IL99/00543	International filing date (day/month/year) 14/10/1999	Priority date (day/month/year) 14/10/1998
International Patent Classification (IPC) or national classification and IPC C12N15/62		
Applicant INTERPHARM LABORATORIES LTD. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.


2. This REPORT consists of a total of 6 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand  03/05/2000	Date of completion of this report  04.12.2000
Name and mailing address of the international preliminary examining authority:   European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer  Wimmer, G  Telephone No. +49 89 2399 7347



# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IL99/00543

## I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).)*:

### Description, pages:

1-21 as originally filed

### Claims, No.:

1-15 as originally filed

### Drawings, sheets:

1/6-6/6 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IL99/00543

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes: Claims 1-15
	No: Claims
Inventive step (IS)	Yes: Claims 2, 6, 8, 15
	No: Claims 1, 3, 5, 7, 9-14
Industrial applicability (IA)	Yes: Claims 1-12
	No: Claims

2. Citations and explanations  
**see separate sheet**

The present application refers to a novel hybrid protein comprising intracellular IL-1 receptor antagonist type II (icIL-1ra-II) and human growth hormone (HGH) signal peptide, expression vectors encoding such fusion proteins, and applications thereof.

**Re Item V**

**Reasoned statement under Art. 35(2) PCT with regard to novelty, inventive step or industrial applicability.**

**The application does not meet the requirements of Art. 33 PCT since claim 1 does not appear to contain an inventive step.**

- 1) Reference is made to the following documents (the document numbering corresponds to their order of citation in the international search report):

D1: WO 96 12022 A (APPLIED RESEARCH SYSTEMS) 25 April 1996 (1996-04-25) cited in the application

D4: PECCEU F. ET AL.: 'Human interleukin 1-beta fused to the human growth hormone signal peptide is N-glycosylated and secreted by Chinese hamster ovary cells.' GENE, vol. 97, 1991, pages 253-258, XP002131134 cited in the application

**Novelty.**

- 2) Document D1 is regarded as being the closest prior art to claim 1. D1 discloses the construction of the icIL-1ra-II gene, and its expression in various cells. However, this document does not describe a fusion of icIL-1ra-II with another protein.

Accordingly, claims referring to DNA constructs encoding such proteins, the proteins themselves, or applications of said DNA constructs or proteins, must be regarded as being novel.

Consequently, claims 1 - 15 are considered to fulfil the requirement for novelty.

Inventive Step.

- 3) The subject-matter of **claim 1** differs from the closest prior art (D1) in that the coding sequence for a signal peptide of a protein, which is normally secreted by human cells, is fused in-frame to the known icIL-1ra-II.

The technical problem was to increase the amount of icIL-1ra-II secreted by the target cell.

This is considered obvious, since secretion signal peptides are routinely fused to a protein of interest to improve secretion.

Claim 1 is therefore not regarded to contain an inventive step.

- 4) For the subject-matter of **claim 2**, D1 can also be regarded as the closest prior art.

Unlike claim 1, the subject-matter of claim 2 specifically contains the Human Growth Hormone signal peptide.

The technical problem again was to increase the amount of icIL-1ra-II secreted by the target cell.

Unlike the employment of a general secretion signal peptide (V.3), the use this specific signal peptide is not regarded as being obvious.

Although the fusion of a protein of interest with the HGH signal peptide, is described in D4, no indications are given to combine the teachings of documents D1 and D4, as D1 does not mention the possible fusion of icIL-1ra-II with another protein, and D4 does not mention the use of the HGH signal peptide in fusion with other proteins than interleukin-1 $\beta$ .

Furthermore, the effect of the fusion of the proteins of D4 is somewhat unclear, as the authors also describe elevated levels of secreted interleukin-1 $\beta$ , upon the mere addition of a Methionine at the N-terminus of the protein.

Consequently, the creation of such a fusion protein is regarded to comprise an

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IL99/00543

inventive step. Accordingly, claim 2 is considered to comply with Art. 33(3) PCT concerning inventivity.

- 5) **Claims 3, 5, 7 and 9 - 14** could only be regarded as to contain an inventive step, if they were based on claim 2, rather than on claim 1.
- 6) **Claims 4, 6, 8 and 15**, which refer back to claim 2, can be regarded as containing an inventive step.

Industrial Applicability.

- 4) For the assessment of the present claims 13 - 15 on the question whether they are industrially applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the treatment of the human or animal body by surgery or therapy, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU.

To:

Assistant Commissioner for Patents  
United States Patent and Trademark  
Office  
Box PCT  
Washington, D.C. 20231  
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

<b>Date of mailing</b> (day/month/year) 24 May 2000 (24.05.00)	
<b>International application No.</b> PCT/IL99/00543	<b>Applicant's or agent's file reference</b> IPL-9 PCT
<b>International filing date</b> (day/month/year) 14 October 1999 (14.10.99)	<b>Priority date</b> (day/month/year) 14 October 1998 (14.10.98)
<b>Applicant</b> AMITAI, Hagit et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

03 May 2000 (03.05.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was  
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer</p> <p>Claudio Borton</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	---

M.H

# INTERNATIONAL COOPERATION TREATY

## PCT

### INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>IPL-9 PCT</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/IL 99/ 00543</b>	International filing date (day/month/year) <b>14/10/1999</b>	(Earliest) Priority Date (day/month/year) <b>14/10/1998</b>
Applicant <b>INTERPHARM LABORATORIES LTD. et al.</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 4 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

#### 1. Basis of the report

- a. With regard to the language, the International search was carried out on the basis of the International application in the language in which it was filed, unless otherwise indicated under this item.

☐ the International search was carried out on the basis of a translation of the International application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any nucleotide and/or amino acid sequence disclosed in the International application, the International search was carried out on the basis of the sequence listing :

☐ contained in the International application in written form.

☐ filed together with the International application in computer readable form.

☒ furnished subsequently to this Authority in written form.

☒ furnished subsequently to this Authority in computer readable form.

☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the International application as filed has been furnished.

☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ Certain claims were found unsearchable (See Box I).

3. ☐ Unity of invention is lacking (see Box II).

#### 4. With regard to the title,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

#### 5. With regard to the abstract,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this International search report, submit comments to this Authority.

#### 6. The figure of the drawings to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☐ None of the figures.



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL 99/00543

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
**Remark: Although claims 13 -15  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IL 99/00543

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/62 C12N15/12 C12N5/10 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 12022 A (APPLIED RESEARCH SYSTEMS) 25 April 1996 (1996-04-25) cited in the application the whole document	1-15
A	----- MUZIO M ET AL: "CLONING AND CHARACTERIZATION OF A NEW ISOFORM OF THE INTERLEUKIN 1 RECEPTOR ANTAGONIST" JOURNAL OF EXPERIMENTAL MEDICINE, JP, TOKYO, vol. 182, no. 2, 1 August 1995 (1995-08-01), pages 623-628, XP000564500 ISSN: 0022-1007 cited in the application the whole document ----- -/--	1-15

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

22 February 2000

Date of mailing of the international search report

09/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gall, I

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IL 99/00543

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HASKILL S. ET AL.: "cDNA cloning of an intracellular form of the human interleukin-1 receptor antagonist associated with the epithelium" PROC. NATL. ACAD. SCI. USA, vol. 88, 1991, pages 3681-3685, XP002131133 cited in the application the whole document	1-15
A	PECCEU F. ET AL.: "Human interleukin 1-beta fused to the human growth hormone signal peptide is N-glycosylated and secreted by Chinese hamster ovary cells." GENE, vol. 97, 1991, pages 253-258, XP002131134 cited in the application the whole document	1-15

# INTERNATIONAL SEARCH REPORT

Info on patent family members

International Application No

P 99/00543

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9612022 A	25-04-1996	IT 1270662 B	07-05-1997
		AU 701471 B	28-01-1999
		AU 3841795 A	06-05-1996
		BR 9509317 A	14-10-1997
		CA 2202470 A	25-04-1996
		EP 0786002 A	30-07-1997
		JP 10509306 T	14-09-1998
		NO 971624 A	30-05-1997
		US 5739282 A	14-04-1998
		US 5837495 A	17-11-1998
		US 5981713 A	09-11-1999

---



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C12N 15/62, 15/12, 5/10, A61K 38/17</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/22146</b> <b>(43) International Publication Date:</b> 20 April 2000 (20.04.00)
<b>(21) International Application Number:</b> PCT/IL99/00543 <b>(22) International Filing Date:</b> 14 October 1999 (14.10.99) <b>(30) Priority Data:</b> 126562 14 October 1998 (14.10.98) IL <b>(71) Applicants (for all designated States except US):</b> INTER-PHARM LABORATORIES LTD. [IL/IL]; Science-Based Industrial Park, Kiryat Weizmann, 76110 Ness-Ziona (IL). APPLIED RESEARCH SYSTEMS ARS HOLDING N.V. [NL/NL]; Pietermaai 15, Curacao (AN). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> AMITAI, Hagit [IL/IL]; Rechov Paldi 4/30, 76248 Rehovot (IL). CHITLARU, Edith [IL/IL]; Hanassi Harishon 16/34, 76302 Rehovot (IL). <b>(74) Agent:</b> EINAV, Henry; Interpharm Laboratories Ltd., Science-based Industrial Park, Kiryat Weizmann, 76110 Ness-Ziona (IL).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> EXPRESSION AND SECRETION OF icIL-1 RECEPTOR ANTAGONIST TYPE II <b>(57) Abstract</b> <p>Novel glycosylated intracellular IL-1 receptor antagonist type II (icIL-1ra-II) is expressed and secreted in mammalian cells transformed with an expression vector where icIL-1ra-II is secreted by expressing icIL-1ra-II fused to the human growth hormone signal peptide. Also disclosed are a pharmaceutical composition containing glycosylated icIL-1ra-II as an active ingredient and a method for reducing IL-1 levels in patients having a condition involving overexpressed IL-1.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## EXPRESSION AND SECRETION OF icIL-1 RECEPTOR ANTAGONIST TYPE II

### Field of the Invention

5           The present invention relates to the expression and secretion of recombinant proteins produced by DNA plasmid expression vectors in mammalian cells. More particularly this invention relates to the recombinant production of intracellular IL-1 receptor antagonist (icIL-1ra) type II by cultured COS and CHO cells, by use of DNA expression vectors containing the genomic DNA sequence of the  
10       human growth hormone (hGH) signal peptide and the cDNA of icIL-1ra type II.

### Background of the Invention

          IL-1 (IL-1 $\alpha$  and IL-1 $\beta$ ) is a pleiotropic cytokine that exerts a variety of effects on different tissues (Dinarello, 1991). IL-1 affects nearly every cell type, either  
15       alone or in synergy with other cytokines (Dinarello, 1996). Two natural pathways of negative regulation strictly control the potent inflammatory effects of IL-1, under physiological conditions. One is IL-1 receptor type II, which is a non-signaling cell-surface IL-1 binding molecule, that acts as a decoy target for IL-1 (Colotta et al, 1993; Sims et al, 1993; Colotta et al, 1994). The second is the unique, IL-1 receptor  
20       antagonist (IL-1ra) (Hannum et al, 1990; Eisenberg et al, 1990; Carter et al, 1990) polypeptide that binds both surface IL-1 receptors, and inhibits signaling from the functional IL-1 receptor.

          Two forms of IL-1ra have been identified. The first was a secreted form, soluble IL-1ra (sIL-1ra), that contains a classical 25-amino acid signal peptide  
25       (Eisenberg et al, 1990; Carter et al, 1990). The second, which does not contain any signal peptide, was termed intracellular IL-1ra (icIL-1ra) (Haskill et al, 1991). icIL-1ra was in fact found to be constitutively expressed intracellularly, in

keratinocytes and in epithelial cells. icIL-1ra was shown to inhibit exogenous IL-1 dependent responses (Haskill et al, 1991).

The two IL-1ra isoforms are derived from the same gene. icIL-1ra transcript originates from an alternative start site, and splicing of an alternative first  
5 exon into an internal splice acceptor site located in the first exon of sIL-1ra (Haskill et al, 1991). These proteins are thus identical, except in their NH<sub>2</sub> end, in which the 21 amino acid signal peptide of sIL-1ra is substituted by three amino acids in icIL-1ra. sIL-1ra and icIL-1ra have a similar capability to inhibit IL-1 activity (Bertini et al, 1992) although expression of the two antagonists is differentially regulated (Haskill et  
10 al, 1991).

An additional isoform of icIL-1ra, termed the type II icIL-1ra, has been recently identified, cloned and functionally characterized (Muzio et al, 1995, WO 96/12022). The type II icIL-1ra contains an additional, in-frame, 63 bp sequence located three codons downstream of the translation start of icIL-1ra. This additional  
15 sequence is inserted between the first and the second exons of the intracellular form of IL-1ra. The additional exon is coded by an extra exon located 2kb downstream of the first icIL-1ra specific exon.

Human growth hormone (hGH) is a 191-amino acid protein synthesized and secreted by the somatotroph cells of the anterior pituitary. The hGH gene contains  
20 five exons and is the best characterized of the five members of the hGH gene family (DeNoto et al, 1981). *In vitro* transfection of the hGH gene into mammalian cells was found to yield high levels of secreted protein proportional to the levels of cytoplasmic hGH mRNA. Thus, secretion does not appear to be the rate-limiting step for appearance of hGH in the culture medium (Selden et al, 1986). The hGH gene  
25 includes a 26 amino acid signal peptide.

Pecceu et al (1991) discloses an attempt to use the human growth hormone signal peptide to create a hybrid gene with the mature form of interleukin-1 $\beta$



(IL-1 $\beta$ ) in order to cause mammalian cells to secrete recombinant IL-1 $\beta$ . Natural IL-1 $\beta$  is expressed initially as an intracellular 31-kDa precursor polypeptide. When proteolytic processing of the precursor occurs, secretion of a mature 17-kDa IL-1 $\beta$  in a soluble mature non-glycosylated form occurs. Pecceu discloses that fusion of the mature form of IL-1 $\beta$  to the heterologous hGH leader sequence permitted the mature IL-1 $\beta$  to be secreted in mature form in CHO cells, although the form which was secreted was a glycosylated form as opposed to the non-glycosylated natural form. Pecceu discloses that the glycosylated form is biologically active. However, Pecceu further states that when the biologically active part of IL-1 $\beta$  was preceded only by a methionine and synthesized in CHO cells, a considerable percentage of the IL-1 $\beta$  produced was quite unexpectedly found in the culture medium. This disclosure leaves some amount of doubt as to whether it was the hGH signal peptide which caused the expression of the IL-1 $\beta$  in the CHO cells or whether such expression was specific to the mechanism involved with this particular protein, the mature form of which is naturally secreted after a precursor protein is expressed intracellularly and then cleaved to form the mature protein which is secreted. Furthermore, Pecceu reports no results as to whether the non-natural glycosylated form of IL-1 $\beta$  creates an immunologic reaction when administered to a human or is recognized as a self protein.

Specific situations involving the recombinant production of non-secretory proteins by fusing a signal peptide of another secretory protein are disclosed in Bjorkdahl et al (1997) and Komada et al (1997).

### **Summary of the Invention**

The present invention provides a method for the production of a recombinant intracellular protein, icIL-1ra type II, in mammalian cells. More particularly, the invention provides a process for engineering proteins to be secreted by use of a signal peptide derived from hGH in different expression vectors and to

produce the secreted proteins in different mammalian cells.

### **Description of the Figures**

Figures 1A-1C show, in Fig. 1A, the genomic hGH signal peptide DNA sequence (SEQ ID NO:1), its amino acid sequence (SEQ ID NO:2), and the primers P1 and P2; in Fig. 1B, the beginning (SEQ ID NO:3) and the end (SEQ ID NO:5) of icIL-1ra type II cDNA, their amino acid sequences (SEQ ID NO:4 and NO:6), and primers P3 and P4 used for construction of the fusion constructs, and in Fig. 1C a schematic representation of templates and primers.

**P1:** hGH-sp 5' primer, containing HindIII restriction site (SEQ ID NO:7).

**P2:** hGH-sp 3' primer, containing 3' icIL-1ra-II sequence overhang (SEQ ID NO:8).

**P3:** icIL-1ra-II 5' primer, containing 5' hGH-sp sequence overhang (SEQ ID NO:9).

**P4:** icIL-1ra-II 3' primer, containing two stop codons and BamHI restriction site (SEQ ID NO:10).

**Figures 2A-2C** describe, in Fig. 2A, the construction of the pCDIC and, in Fig. 2B, the construction of pSGHIRA2 DNA vectors used for expression of the icIL-1ra type II in mammalian cells. Fig. 2C is a scheme of pDHFR.

### **Detailed Description of the Invention**

The natural form of icIL-1ra type II is expressed intracellularly and is not secreted by the cells in which it is produced. However, in accordance with the present invention, this protein can be secreted in a mammalian recombinant production system by fusing the DNA encoding the protein to the DNA encoding the signal peptide of another human protein which is normally expressed and secreted by human

cells, and which is known to cause expression of the human protein in non-human mammalian expression systems. Preferably, the signal peptide is the 26-amino acid signal peptide of the human growth hormone gene.

Prior to the present invention, it could not have been reasonably  
5 predicted whether or not the hGH signal peptide would drive the expression of icIL-1ra-II in a mammalian cell expression system in view of the fact that icIL-1ra-II is naturally expressed only intracellularly and is not secreted from the cell. In the known prior art, as represented by Pecceu et al (1991), the hGH signal peptide was used to express and secrete the mature form of IL-1 $\beta$ . However, the mature form of  
10 IL-1 $\beta$  is naturally secreted from the cells in which it is produced, although indirectly. A precursor protein is first produced which, after intracellular processing, is secreted from the cell. However, Pecceu discloses that when a recombinant vector containing only the DNA encoding the mature form of IL-1 $\beta$ , without any signal protein, is used, the protein is secreted from CHO cells. Thus, it could not be predicted with a  
15 reasonable degree of certainty that a protein such as icIL-1ra-II, which is only expressed intracellularly and is not naturally secreted from the cell, could be made to be secreted in large quantities in a recombinant mammalian expression system when fused to an hGH signal peptide or to a signal peptide of another secretory protein.

The icIL-1ra-II protein produced in accordance with the present  
20 invention is glycosylated while the natural protein is non-glycosylated. Thus, the present invention further relates to the two novel glycosylated forms of icIL-1ra-II produced for the first time by means of the present invention. These are the glycosylated forms which have apparent molecular weight of approximately 27 kDa and 30 kDa as determined by Commassie blue staining of SDS-PAGE (15%  
25 acrylamide under reducing conditions). It could not be predicted with a reasonable degree of certainty whether these novel glycosylated forms of icIL-1ra-II will retain the biological activity of natural icIL-1ra-II and will not be immunogenic when

administered to humans. Experiments with these two novel glycosylated forms of icIL-1ra-II will establish that they are indeed biologically active and non-immunogenic when administered to humans.

Accordingly, the present invention is directed to a process for the  
5 recombinant expression of a protein having the amino acid sequence of natural icIL-1ra-II in a recombinant cell expression system through use of a vector which is a fusion of the signal peptide of a human secretory protein, preferably the 26 amino acid signal peptide of hGH; fused in proper reading frame with the DNA encoding icIL-1ra-II. The process comprises producing an expression vector containing DNA  
10 encoding icIL-1ra-II, either in the form of cDNA or genomic DNA, fused in proper reading frame with DNA encoding the selected signal peptide, preferably the 26 amino acid hGH signal peptide. The expression vector is then inserted into an appropriate expression host, such as CHO cells. The transformed host cells are then cultured in a manner which causes the expression vector to express its encoded protein and the  
15 expressed and secreted icIL-1ra-II protein is then collected and purified from the culture medium.

The present invention is not intended to be limited by the specific examples presented herein. While CHO cells are used as the host cells, any other eukaryotic expression system, preferably mammalian expression system, may be used  
20 such as COS cells, yeast cells, insect cells, etc. Those of ordinary skill in the art are well aware of the techniques of creating expression vectors, inserting them into expression systems and selecting clones which express the desired protein, including amplification techniques.

As would be appreciated by those skilled in the art, the types of  
25 promoters used to control transcription of the icIL-ra-II proteins may be any of those which are functional in the host cells. Examples of promoters functional in mammalian cells include the SV40 early promoter, adenovirus major late promoter,

herpes simplex (HSV) thymidine kinase promoter, rous sarcoma (RSV) LTR promoter, human cytomegalovirus (CMV) immediate early promoter, mouse mammary tumor virus (MMTV) LTR promoter, interferon- $\beta$  promoter, heat shock protein 70 (hsp 70) promoter, as well as many others well known in the art. These promoters  
5 may be either constitutive or regulatable. All else being equal, constitutive promoters are preferred because an extra treatment step, such as temperature shift, addition of chemical agents or inducers, etc., is not required for expression from constitutive promoters.

10 The technical advance of the present invention lies in the confirmation that icIL-1ra-II can be secreted in such an expression system when using a signal peptide of a human secretory protein, preferably hGH. All of the other techniques involved are well known to those of ordinary skill in this art and can be practiced without undue experimentation using only the knowledge of the skill of the art available at the time of the present invention.

15 The present invention further is directed to the expression vector which contains the icIL-1ra-II DNA fused to the DNA encoding a signal protein of a human secretory protein, such as hGH, and host cells transfected with such an expression vector.

20 The present invention is further directed to the novel glycosylated forms of icIL-1ra-II produced in accordance with the present invention.

The invention further relates to methods for reducing the amount of IL-1 in patients having a condition involving the overexpression of IL-1, by administering one of the novel glycosylated icIL-1ra-II proteins in accordance with the present invention in a therapeutically effective amount. Appropriate therapeutic dosages for  
25 the reduction of IL-1 in patients having such a condition, can be readily empirically determined by those of ordinary skill in the art.

The glycosylated icIL-1ra-II proteins of the present invention may be

administered by any means that achieves its intended purpose. For example, administration may be by a number of different parenteral routes including, but not limited to, subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intracerebral, intranasal, oral, transdermal, or buccal routes. Parenteral administration  
5 can be bolus injection or by gradual perfusion over time.

It is understood that the dosage administered will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. The total dose required for each treatment may be administered by multiple doses or in a single dose. By  
10 "effective amount", it is meant a concentration of glycosylated icIL-1ra-II protein which is capable of reducing the amount of IL-1 in patients having a condition involving elevated levels of IL-1. Such concentrations can be routinely determined by those of skill in the art. It will also be appreciated by those of skill in the art that the dosage may be dependent on the stability of the administered protein. A less stable  
15 protein may require administration in multiple doses.

The invention also relates to pharmaceutical compositions comprising the glycosylated icIL-1ra-II protein of the present invention with a pharmaceutically acceptable excipient.

Preparations for parenteral administration include sterile aqueous or  
20 non-aqueous solutions, suspensions, and emulsions, which may contain auxiliary agents or excipients which are known in the art.

Pharmaceutical compositions comprising the glycosylated icIL-1ra-II protein according to the invention include all compositions wherein the protein is contained in an amount effective to achieve its intended purpose. In addition, the  
25 pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Suitable

pharmaceutically acceptable vehicles are well known in the art and are described for example in Gennaro, Alfonso, Ed., Remington's Pharmaceutical Sciences, 18th Edition, Mack Publishing Co., Easton, PA (1990), a standard reference text in this field. Pharmaceutically acceptable vehicles can be routinely selected in accordance with the mode of administration and the solubility and stability of the protein. For example, formulations for intravenous administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

While any excipients known for the administration of therapeutic proteins can be used in accordance with the present invention, excipients used for intravenous administration are preferred.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts. In addition, suspension of the active compound as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions that may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

The vectors in accordance with the present invention can also be used in gene therapy to cause appropriate human cells to express icIL-1ra-II *in vivo* in order to direct the IL-1 antagonizing effect of this protein directly at the desired site. The novelty of this process lies in the particular vector and the knowledge of the activity of the glycosylated protein produced thereby and not in the specific methods of gene therapy, including the methods of introducing an expressible vector directly into the cells of interest. These are within the skill of those of ordinary skill in this art at the time the present invention was made.

Non-limiting specific examples of the present invention follow.

**Example 1: Generation of hGH-sp-icIL-1ra-II Fragment**

The human growth hormone hGH signal peptide sequence was amplified  
5 by PCR using the pXGH5 vector, which encodes the full-length human growth  
hormone genomic sequence, as template (DeNoto et al, 1981). pXGH5 was used as a  
template for PCR using primers P1 (containing a HindIII restriction site and a Kozak  
sequence) and P2 (which has a 3' icIL-1ra-II sequence overhang, Fig. 1A). The  
icIL-1ra-II cDNA was amplified by PCR with primers P3 (which has a 5' hGH signal  
10 peptide sequence overhang) and P4 (which has two stop codons and a BamHI  
restriction site, Fig. 1B). These two PCR fragments were annealed together by their  
homologous regions and were further amplified by PCR using primers P1 and P4, to  
generate the hGH-icIL-1ra-II fragment (Fig. 1C).

**Example 2: Construction of pCDIC**

hGH-sp-icIL-1ra-II fragment was digested with HindIII and BamHI and  
cloned into the HindIII-BamHI sites of pCDNA3.1 (+) (Invitrogen, San Diego, Fig.  
2A) downstream of the CMV promoter. The resulting vector (pCDIC, Fig. 2A) was  
mapped by restriction analysis, and used to transfect COS cells.

20

**Example 3: Construction of pSGHIRA2**

hGH-sp-icIL-1ra-II fragment was digested with HindIII and BamHI, and  
cloned into HindIII-BclI sites of pSVE3 (Fig. 2B; Hartman et al, 1982), downstream  
of the early SV-40 promoter. The resulting vector pSGHIRA2 (Fig. 2B) was used to  
transfect CHO DUKX (ATCC, CRL 9010) cells in co-transfection with the mouse  
25 DHFR containing vector pDHFR (Fig. 2C) as detailed below. The constructs were  
analyzed by restriction mapping and sequenced.



**Example 4: Expression Vector Carrying the Mouse DHFR Gene**

Plasmid pDHFR (Fig. 2C) is composed of the complete pBR322 sequence, the SV40 early gene promoter, the 70 bp splicing region of the mouse  $\gamma$ 2a gene fused to the mouse DHFR cDNA, followed by the SV40 early gene polyadenylation signal.

**Example 5: Transient Expression in COS Cells**

pCDIC DNA was used for transfecting COS cells by means of the DEAE dextran method. Cells were seeded at approximately  $3 \times 10^6/80 \text{ cm}^2$  flask and allowed to grow overnight. The next day all medium was aspirated from the flasks and 5 ml of transfection medium was added to the cells. Transfection medium contains 400  $\mu\text{g/ml}$  DEAE dextran 100  $\mu\text{M}$  Chloroquine, 2  $\mu\text{g/ml}$  DNA, and 10% NuSerum in RPMI medium. After incubation for 3-4 hrs at 37°C the transfection medium was removed by aspiration and replaced with 5 ml of 10% DMSO in PBS for 2 minutes at room temperature. This solution was then aspirated and culture medium containing 10% FBS in RPMI added to the flasks. The cultures were incubated at 37°C for 24 hours, then the culture medium was changed to medium containing 2% serum. 24 hours later incubation temperature was reduced to 32°C, and culture supernatant samples were analyzed for the presence of the icIL-1ra type II in the culture supernatant by ELISA (see Example 7). 6-7  $\mu\text{g}/1 \times 10^6$  cells/run icIL-1ra type II were secreted from the transfected COS cells. Highest production levels were at days 4 to 8 following transfection. These results indicate that fusion of the hGH signal peptide to the intracellular form of IL-1-1ra type II, enables its secretion into the culture medium of the transfected cells.

**Example 6: Stable Expression in CHO Cells**

CHO cells were cotransfected with pSGHIRA2 vector carrying the genes for icIL-1ra type II and with pDHFR carrying the gene for mouse DHFR (described in Fig. 2C), by means of the Lipofectamine transfection method.

5                Cells were seeded  $1 \times 10^6$ /10 cm plate in F12 medium containing 10% FCS, and allowed to grow overnight. The cells were washed in F12 medium, and 8 ml of the DNA-Lipofectamine mixture was added to the plates, which were then incubated for 4-5 hours at 37°C. At the end of the incubation period, 8 ml of F12 medium containing 20% FCS were added, and the plates were incubated for 24 hours  
10                at 37°C. The culture medium was then changed to fresh F12 medium containing 10% FCS. 72 hours following transfection cell cultures were seeded, either by limiting dilution, or by subculture at a 1:20 dilution, into selective medium depleted of Thymidine and Hypoxanthine, containing 10% dialyzed FCS, and allowed to grow until  
15                single colonies could be picked and analyzed. Expression of icIL-1ra-II by eight stably integrated CHO clones is summarized in Table 1. The process lends itself to scale up by methods known in the art.

**Table 1****Specific Productivity of icIL-1ra-II by CHO Clones**

5

Clone Number	Specific Productivity ng/10 <sup>6</sup> cells/day	
	Before MTX Amplification	After MTX Amplification (up to 400 nM MTX)
1-33	118	273
2-56	91	220
1-64	122	186
1-84	194	245
2-2	92	524
2-66	124	283
2-73	82	218
2-88	120	442

10

The MTX amplification was performed as follows:

Cells that grew in the absence of MTX were seeded to six T-flasks, in the presence of different MTX concentrations (e.g., 0, 2nM, 5nM, 10nM, 20nM, 50nM). About 10 days later, the cultures were observed microscopically and the cells were counted, in order to determine survival. The MTX concentration that allowed the survival of approximately 10% of the culture was selected for further propagation.

The second round of amplification was performed in a similar manner, however the MTX concentrations were higher, starting from the MTX concentration that was selected at the first round. The cultures were again scored for survival, relative to the control MTX concentration of this round. The clones presented in Table 1 were amplified to the MTX concentrations shown in Table 2.

**Table 2**

Clone	MTX First Round	MTX Second Round
1-33	20nM	300nM
1-64	20nM	400nM
1-84	20nM	100nM
2-2	50nM	200nM
2-56	50nM	100nM
2-66	20nM	100nM
2-73	20nM	100nM
2-88	20nM	100nM

**Example 7: ELISA Test**

Microtiter plates (Nunc) were coated with mouse anti IL-1ra antibody (purified ascitis IgG, MCA 1467, clone 1384, Serotec Ltd, Oxford, UK) 5 µg/ml in  
5 PBS (100 µl/well), for 3 hrs at 37°C, and stored 40°C. The plates were washed with PBS containing Tween 20 (0.05%, referred to herein as washing buffer) and blocked with the same solution containing 1% bovine serum albumin (BSA, referred to herein as blocking solution) for 1 hour at 37°C. Plates were then washed in washing buffer. The samples to be analyzed were diluted in the blocking solution, and added to the  
10 wells (100 µl/well) for 90 minutes at 37°C. The plates were then washed 6 times in washing buffer, followed by addition of biotinylated anti human IL-1ra antibody (100 µl/well of a 1:10,000 dilution, MCA 1466B, clone 1390m, Serotec, Oxford, UK). Plates were incubated for 1 hr at 37°C and washed with washing buffer. A horseradish peroxidase (HRP) streptavidin conjugate (1 mg/ml Sigma, Rehovot, Israel,  
15 100 µl/well diluted 1:10,000 in blocking buffer) was then added to the plates, and incubated for 1.5 hours at 37°C. The plates were then washed in buffer and the substrate solution (*o*-phenylenediamine dihydrochloride, OPD, Sigma Rehovot, Israel, 100 µl/well) was added for 10 min. at 22°. The reaction was stopped by addition of 100 µl/well of 4N HCl. The plates were then read in an automated Elisa reader. A  
20 standard preparation of IL-1ra (Serotec, Oxford, UK, PHP080, 2-128 ng/ml) was used as reference for the IL-1ra concentration.

**Example 8: Affinity Chromatography of icIL-1ra Type II with Monoclonal Antibodies**

25 Affinity chromatography of icIL-1ra type II was performed by binding anti-human IL-1ra antibodies (purified ascitis IgG, MCA 1467, clone 1384, Serotec Ltd, Oxford, UK) to CNBr activated Sepharose 4B (5 mg/ml resin, Pharmacia,

Uppsala, Sweden). Culture supernatant from CHO cells, of clone 2-88 of the above-mentioned Example 6 was diafiltrated over a 100K membrane and then concentrated over a 10K membrane. Concentrated proteins were dialyzed against 0.1 M  $\text{NaHCO}_3$ , 150 mM NaCl pH 8.2. This procedure enriched the product concentration, reduced the volume of sample (100 fold) and removes major impurities. The yield of this step is about 85%. 30 ml of concentrated proteins were loaded on a 3.2 ml column, that had been equilibrated with 0.1 M sodium carbonate, 150 mM sodium chloride, pH 8.3, at a flow rate of 2 ml/min. icIL-1ra was eluted in 150 mM citric acid, 300 mM NaCl pH 2.7. Eluted fractions were immediately neutralized with 1M Tris pH 9.3. The fraction eluted resolved into two bands of apparent molecular weight of approximately 27 kDa and 30 kDa respectively, as determined by Commassie blue staining of SDS-PAGE (15% acrylamide under reducing conditions). The different molecular weights are presumably due to variations in glycosylation.

#### **Example 9: Western Blot**

The affinity chromatography column eluate of above-mentioned Example 8 was concentrated by filtration on a 3K membrane (Miniset, Pall Filtron, Northborough, MA). Fractions were resolved on a 15% acrylamide SDS-PAGE gel under reducing conditions (Readygel BioRad, Hercules, CA) and electroblotted onto a nitrocellulose membrane (BRL, Life Technologies, MD). The blot was incubated in PBS containing 10% low fat milk, 0.1% Tween 20, overnight. The blot was then incubated with mouse anti human IL-1ra antibodies (purified ascitis IgG 1:5,000, MCA 1467, Serotec Ltd, Oxford, UK) for 2 hours at RT, then washed three times for 15 minutes in PBS containing 0.1% Tween 20, and further incubated with goat anti-mouse horseradish peroxidase-alkaline phosphatase (1:10,000 Sigma, Israel) for 1 hour at RT. The blot was then washed 3 times in PBS containing 0.1% Tween 20, followed by detection with enhanced luminescence (Amersham). Two protein bands

of approximately 27 kDa and 30 kDa, respectively, corresponding to icIL-1ra, were identified.

#### **Example 10: Protein Sequence Analysis**

5           The purified fraction from the immunoaffinity chromatography column of Example 8 was electroblotted in parallel, both on a PVDF membrane (Millipore, Bedford, MA), and on a nitrocellulose membrane for Western blotting analysis as described in the above-mentioned Example 9. The two bands stained by Coomassie blue were both recognized as IL-1ra by Western blot analysis. The purified fraction  
10           eluted from the affinity chromatography column, as well as the two bands excised from the Coomassie blue stained PVDF membrane, were subjected to protein sequence analysis by Edman degradation in the Procise™, 491HT microsequencer (Applied Biosystems, USA). Sequencing of the N-terminal amino acids, indicated that the purified fraction separated from the culture supernatant contained two forms of  
15           icIL-1ra. The amino acid sequence obtained, ALADLYEEGGGGGGE (SEQ ID NO:11), demonstrated that the secreted protein represented the mature icIL-1ra type II protein beginning at amino acid position +2 from the deduced start of translation of the gene (GENBANK, ID# X84348). An additional icIL-1ra form, beginning at amino acid +1 from the deduced start of translation of the icIL-1ra type II, was found  
20           as well.

          Having now fully described this invention, it will be appreciated by those skilled in the art that the same can be performed within a wide range of equivalent parameters, concentrations, and conditions without departing from the spirit and scope of the invention and without undue experimentation.

25           While this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the

inventions following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth as follows in the scope of the appended claims.

5 All references cited herein, including journal articles or abstracts, published or unpublished U.S. or foreign patent applications, issued U.S. or foreign patents, or any other references, are entirely incorporated by reference herein, including all data, tables, figures, and text presented in the cited references. Additionally, the entire contents of the references cited within the references cited  
10 herein are also entirely incorporated by reference.

Reference to known method steps, conventional method steps, known methods or conventional methods is not in any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

15 The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art (including the contents of the references cited herein), readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present  
20 invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the  
25 skilled artisan in light of the teachings and guidance presented herein, in combination with the knowledge of one of ordinary skill in the art.



**REFERENCES**

- Bertini et al, "Inhibitory effect of recombinant intracellular interleukin 1 receptor antagonist on endothelial cell activation", Cytokine 4(1):44-47 (1992)
- 5 Bjorkdahl et al, "Gene transfer of a hybrid interleukin-1 beta gene to B16 mouse melanoma recruits leucocyte subsets and reduces tumour growth in vivo", Cancer Immunol Immunother 44(5):273-81 (1997)
- 10 Carter et al, "Purification, cloning, expression and biological characterization of an interleukin-1 receptor antagonist protein", Nature 344(6267):633-638 (1990)
- Colotta et al, "Interleukin-1 type II receptor: a decoy target for IL-1 that is regulated by IL-4", Science 261(5120):472-475 (1993)
- 15 Colotta F et al, "The type II 'decoy' receptor: a novel regulatory pathway for interleukin 1, Immunol Today 15(12):562-566 (1994)
- 20 DeNoto et al, "GoHM Human growth hormone DNA sequence and mRNA structure: possible alternative splicing", Nucleic Acids Res 9(15):3719-3730 (1981)
- Dinarelli, CA "Interleukin-1 and interleukin-1 antagonism", Blood 77(8):1627-1652 (1991)
- 25 Dinarelli, CA "Biologic basis for interleukin-1 in disease", Blood 87(6):2095-2147 (1996)
- Eisenberg et al, "Primary structure and functional expression from complementary

DNA of a human interleukin-1 receptor antagonist", Nature 343(6256):341-346 (1990)

Gennaro, Alfonso, Ed., Remington's Pharmaceutical Sciences, 18th Edition 1990,  
5 Mack Publishing Co., Easton, PA,

Hannum et al, "Interleukin-1 receptor antagonist activity of a human interleukin-1 inhibitor", Nature 343(6256):336-340 (1990)

10 Hartman et al, "Human influenza virus hemagglutinin is expressed in monkey cells using simian virus 40 vectors", Proc Natl Acad Sci USA, 79 (2):233-237 (1982)

Haskill et al, "cDNA cloning of an intracellular form of the human interleukin 1 receptor antagonist associated with epithelium, Proc Natl Acad Sci USA  
15 88(9):3681-3685 (1991)

Komada et al, "Protective effect of transfection with secretable superoxide dismutase (SOD) (a signal sequence-SOD fusion protein coding cDNA) expression vector on superoxide anion-induced cytotoxicity in vitro", Biol Pharm Bull  
20(5):530-6 (1997)

20

Muzio et al, "Cloning and characterization of a new isoform of the interleukin 1 receptor antagonist," J Exp Med 182(2):623-628 (1995)

Pecceu et al, "Human interleukin 1 $\beta$  fused to the human growth hormone signal  
25 peptide is N-glycosylated and secreted by Chinese hamster ovary cells", Gene 97(2):253-258 (1991)

Selden et al, "Human growth hormone as a reporter gene in regulation studies employing transient gene expression", Mol Cell Biol 6(9):3173-3179 (1986)

5 Sims et al, "Interleukin 1 signaling occurs exclusively via the type I receptor", Proc Natl Acad Sci USA, 90(13):6155-6159 (1993)

**CLAIMS**

1. An expression vector, comprising a DNA segment encoding a signal peptide of a protein which is normally expressed and secreted by human cells, joined to a DNA segment encoding intracellular IL-1 receptor antagonist type II (icIL-1ra-II) and operably linked to a promoter sequence, wherein said icIL-1ra-II is expressed from said promoter sequence and translated with said signal peptide fused in frame to icIL-1ra-II.
2. An expression vector in accordance with claim 1, wherein said signal peptide is human growth hormone signal peptide.
3. A host cell transformed with the expression vector of claim 1.
4. A host cell transformed with the expression vector of claim 2.
5. A host cell in accordance with claim 3, wherein said cell is an endogenous cell of a human host.
6. A host cell in accordance with claim 4, wherein said cell is an endogenous cell of a human host.
7. A method for producing a recombinant icIL-1ra-II comprising the steps of:
  - culturing a host cell according to claim 3 to express and produce a recombinant glycosylated icIL-1ra-II;
  - recovering the produced recombinant glycosylated icIL-1ra-II.
8. A method for producing a recombinant icIL-1ra-II comprising the steps of:
  - culturing a host cell according to claim 4 to express and produce a recombinant glycosylated icIL-1ra-II;
  - recovering the produced recombinant glycosylated icIL-1ra-II.
9. A glycosylated icIL-1ra-II produced by a method according to claim 7.

10. The glycosylated icIL-1ra-II according to claim 9 having an apparent molecular weight of about 27 kDa on SDS-PAGE under reducing conditions with 15% acrylamide.

5 11. The glycosylated icIL-1ra-II according to claim 9 having an apparent molecular weight of about 30 kDa on SDS-PAGE under reducing conditions with 15% acrylamide.

12. A pharmaceutical composition, comprising the glycosylated icIL-1ra-II according to claim 9 in a therapeutically effective amount and a pharmaceutically acceptable excipient.

10 13. A method for reducing the amount of IL-1 in a patient having a condition associated with overexpression of IL-1, comprising administering the pharmaceutical composition according to claim 12 to a patient in need thereof.

14. A method for reducing the amount of IL-1 at a desired site in a human patient, comprising introducing a vector in accordance with claim 3 into  
15 appropriate endogenous human cells at the desired site to produce transformed cells which will express icIL-1ra-II at the desired site.

15. A method for reducing the amount of IL-1 at a desired site in a human patient, comprising introducing a vector in accordance with claim 4 into  
20 appropriate endogenous human cells at the desired site to produce transformed cells which will express icIL-1ra-II at the desired site.

A Human GH Signal Peptide Genomic Sequence

5' ————— 3' →  
P1 CCCCAAGCTTGCCACC ATG GCT ACA G GTAAGCGC  
HindIII  
ATG GCT ACA G GTAAGCGCCC CTAAATCCC TTGGGCACA  
Met Ala Thr G



ATGTGTCTG AGGGAGAGG CAGGACCTG TAGATGGGAC GGGGGCACTA ACCCTCAGGT  
TTGGGGCTTC TGAATGTGAG TATCGCCATG TAAGCCCAGT ATTTGGCCAA TCTCAGAAAG  
CTCCTGGTCC CTGGAGGGAT GGAGAGAGAA AAACAAACAG CTCCTGGAGC AGGGAGAGTG  
CTGGCCTCTT GCTCTCCGGC TCCCTCTGTT GCCCTCTGGT TTCTCCCCAG

GC TCC CGG ACG TCC CTG CTC CTG GCT TTT GGC CTG CTC TGC  
ly Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys

3' ← ————— 5'  
P2 C TGG CTT CAA GAG GGC AGT GCC ATG GCT TTA GCT GAC  
CTG CCC TGG CTT CAA GAG GGC AGT GCC  
Leu Pro Trp Leu Gln Glu Gly Ser Ala  
↑  
signal peptide cleavage site

Fig. 1A

2/6

**B****icIL1RaII, cDNA sequence**

BamHI

GTC ACC AAA TTC TAC TTC CAG GAG GAC GAG TAG

Val Thr Lys Phe Tyr Phe Gln Glu Asp Glu ---

Primers are boxed, direction of synthesis is indicated by arrows, and overhangs are bold.

Enzymatic restriction sites are underlined.

Fig. 1B

3/6




Fig. 1C



4/6




Fig. 2A




Fig. 2B

6/6




Fig. 2C

# INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/IL 99/00543

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/62 C12N15/12 C12N5/10 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 12022 A (APPLIED RESEARCH SYSTEMS) 25 April 1996 (1996-04-25) cited in the application the whole document ---	1-15
A	MUZIO M ET AL: "CLONING AND CHARACTERIZATION OF A NEW ISOFORM OF THE INTERLEUKIN 1 RECEPTOR ANTAGONIST" JOURNAL OF EXPERIMENTAL MEDICINE, JP, TOKYO, vol. 182, no. 2, 1 August 1995 (1995-08-01), pages 623-628, XP000564500 ISSN: 0022-1007 cited in the application the whole document --- -/--	1-15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 February 2000

Date of mailing of the international search report

09/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

# INTERNATIONAL SEARCH REPORT

Int. Patent Application No.

PCT/IL 99/00543

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HASKILL S. ET AL.: "cDNA cloning of an intracellular form of the human interleukin-1 receptor antagonist associated with the epithelium" PROC. NATL. ACAD. SCI. USA, vol. 88, 1991, pages 3681-3685, XP002131133 cited in the application the whole document ---	1-15
A	PECCEU F. ET AL.: "Human interleukin 1-beta fused to the human growth hormone signal peptide is N-glycosylated and secreted by Chinese hamster ovary cells." GENE, vol. 97, 1991, pages 253-258, XP002131134 cited in the application the whole document -----	1-15

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL 99/00543

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 13 -15  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 99/00543

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9612022 A	25-04-1996	IT 1270662 B	07-05-1997
		AU 701471 B	28-01-1999
		AU 3841795 A	06-05-1996
		BR 9509317 A	14-10-1997
		CA 2202470 A	25-04-1996
		EP 0786002 A	30-07-1997
		JP 10509306 T	14-09-1998
		NO 971624 A	30-05-1997
		US 5739282 A	14-04-1998
		US 5837495 A	17-11-1998
		US 5981713 A	09-11-1999

# PATENT COOPERATION TREATY

From the  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

## PCT

To:

EINAV, Henry  
INTERPHARM LABORATORIES LTD.  
Science-based Industrial Park  
Kiryat Weizmann  
Ness-Ziona 76110  
ISRAEL

אינטרפרם מעבדות בע"מ  
17. 12. 2000  
דניאל

NOTIFICATION OF TRANSMITTAL OF  
THE INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT  
(PCT Rule 71.1)

Date of mailing  
(day/month/year) 04.12.2000

Applicant's or agent's reference  
? 186-9

### IMPORTANT NOTIFICATION

International application No.  
PCT/IL99/00543

International filing date (day/month/year)  
14/10/1999

Priority date (day/month/year)  
14/10/1998

Applicant  
INTERPHARM LABORATORIES LTD. et al.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

#### 4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/



European Patent Office  
D-80298 Munich  
Tel. +49 89 2399 - 0 Tx: 523656 epmu d  
Fax: +49 89 2399 - 4465

Authorized officer

Emslander, S

Tel. +49 89 2399-8718





# PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference ?	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/IL99/00543	International filing date (day/month/year) 14/10/1999	Priority date (day/month/year) 14/10/1998
International Patent Classification (IPC) or national classification and IPC C12N15/62		
Applicant INTERPHARM LABORATORIES LTD. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 6 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand  03/05/2000	Date of completion of this report  04.12.2000
Name and mailing address of the international preliminary examining authority:   European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer  Wimmer, G  Telephone No. +49 89 2399 7347 

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IL99/00543

## I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

**Description, pages:**

1-21 as originally filed

**Claims, No.:**

1-15 as originally filed

**Drawings, sheets:**

1/6-6/6 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IL99/00543

☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims	1-15
	No:	Claims	
Inventive step (IS)	Yes:	Claims	2, 6, 8, 15
	No:	Claims	1, 3, 5, 7, 9-14
Industrial applicability (IA)	Yes:	Claims	1-12
	No:	Claims	

### 2. Citations and explanations see separate sheet

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IL99/00543

The present application refers to a novel hybrid protein comprising intracellular IL-1 receptor antagonist type II (icIL-1ra-II) and human growth hormone (HGH) signal peptide, expression vectors encoding such fusion proteins, and applications thereof.

**Re Item V**

**Reasoned statement under Art. 35(2) PCT with regard to novelty, inventive step or industrial applicability.**

**The application does not meet the requirements of Art. 33 PCT since claim 1 does not appear to contain an inventive step.**

- 1) Reference is made to the following documents (the document numbering corresponds to their order of citation in the international search report):

D1: WO 96 12022 A (APPLIED RESEARCH SYSTEMS) 25 April 1996 (1996-04-25) cited in the application

D4: PECCEU F. ET AL.: 'Human interleukin 1-beta fused to the human growth hormone signal peptide is N-glycosylated and secreted by Chinese hamster ovary cells.' GENE, vol. 97, 1991, pages 253-258, XP002131134 cited in the application

**Novelty.**

- 2) Document D1 is regarded as being the closest prior art to claim 1. D1 discloses the construction of the icIL-1ra-II gene, and its expression in various cells. However, this document does not describe a fusion of icIL-1ra-II with another protein.

Accordingly, claims referring to DNA constructs encoding such proteins, the proteins themselves, or applications of said DNA constructs or proteins, must be regarded as being novel.

Consequently, claims 1 - 15 are considered to fulfil the requirement for novelty.

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IL99/00543

Inventive Step.

- 3) The subject-matter of **claim 1** differs from the closest prior art (D1) in that the coding sequence for a signal peptide of a protein, which is normally secreted by human cells, is fused in-frame to the known icIL-1ra-II.

The technical problem was to increase the amount of icIL-1ra-II secreted by the target cell.

This is considered obvious, since secretion signal peptides are routinely fused to a protein of interest to improve secretion.

Claim 1 is therefore not regarded to contain an inventive step.

- 4) For the subject-matter of **claim 2**, D1 can also be regarded as the closest prior art.

Unlike claim 1, the subject-matter of claim 2 specifically contains the Human Growth Hormone signal peptide.

The technical problem again was to increase the amount of icIL-1ra-II secreted by the target cell.

Unlike the employment of a general secretion signal peptide (V.3), the use this specific signal peptide is not regarded as being obvious.

Although the fusion of a protein of interest with the HGH signal peptide, is described in D4, no indications are given to combine the teachings of documents D1 and D4, as D1 does not mention the possible fusion of icIL-1ra-II with another protein, and D4 does not mention the use of the HGH signal peptide in fusion with other proteins than interleukin-1 $\beta$ .

Furthermore, the effect of the fusion of the proteins of D4 is somewhat unclear, as the authors also describe elevated levels of secreted interleukin-1 $\beta$ , upon the mere addition of a Methionine at the N-terminus of the protein.

Consequently, the creation of such a fusion protein is regarded to comprise an

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IL99/00543

inventive step. Accordingly, claim 2 is considered to comply with Art. 33(3) PCT concerning inventivity.

- 5) **Claims 3, 5, 7 and 9 - 14** could only be regarded as to contain an inventive step, if they were based on claim 2, rather than on claim 1.
- 6) **Claims 4, 6, 8 and 15**, which refer back to claim 2, can be regarded as containing an inventive step.

**Industrial Applicability.**

- 4) For the assessment of the present claims 13 - 15 on the question whether they are industrially applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the treatment of the human or animal body by surgery or therapy, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.